WHAT IS CLAIMED IS:

1.	A medical	device	comprising

an implantable structure;

5

a basecoat matrix, including a combination of rapamycin and 2-methoxyestradiol, in therapeutic dosages, incorporated in a first polymeric material, the basecoat matrix being affixed to the surface of the implantable medical device; and

10

a topcoat, including a second polymeric material, affixed to the basecoat matrix for controlling the elution rate of the rapamycin and the 2-methoxyestradiol.

2. The medical device according to claim 1, wherein the implantable structure comprises a stent.

15

3. The medical device according to claim 1, wherein the implantable structure comprises a stent-graft.

20

4. The medical device according to claim 1, wherein the implantable structure comprises an anastomosis device.

5. The medical device according to claim 1, wherein the second polymeric material is incompatible with the first polymeric material, thereby creating both a physical and chemical barrier to the elution of the rapamycin and the 2-methoxyestradiol.

25

6. The medical device according to claim 5, wherein the first polymeric material comprises a fluoropolymer.

30

7. The medical device according to claim 6, wherein the second polymeric material comprises an acrylic.

5

15

8.	A medical device comprising:
	an implantable structure; and

- a combination of rapamycin and 2-methoxyestradiol, in therapeutic dosages, releasably affixed to the implantable structure for the treatment of restenosis following vascular injury.
- 9. The medical device according to claim 8, wherein the implantable structure comprises a stent.
- 10. The medical device according to claim 8, wherein the implantable structure comprises a stent-graft.
 - 11. The medical device according to claim 8, wherein the implantable structure comprises an anastomosis device
 - 12. The medical device according to claim 8, further comprising a polymeric coating, the combination of rapamycin and 2-methoxyestradiol being incorporated into the polymeric coating.
- 20 13. A method for treating restenosis comprising the local administration of a therapeutic dose of a combination of rapamycin and 2-methoxyestradiol.
- 14. A method for treating restenosis comprising the administration of a therapeutic dose of a combination of rapamycin and 2-methoxyestradiol.